

southampton dataobservatory

Joint Strategic Needs Assessment (JSNA) August Review

Southampton City Council



- Health & Wellbeing Boards are responsible for producing a JSNA (Health & Social Care Act 2012)
- The JSNA is an assessment of the current and future health and social care needs of the community
- Purpose is to improve health & wellbeing and reduce inequalities
- Statutory requirement to produce AND inform health and wellbeing commissioning plans
- Locally determined process No mandated format, core dataset or update schedule. Southampton
 JSNA is brought together with other data, intelligence, specialist reports, needs assessments,
 summary analysis and headline statistics covering the city's population, health, community safety,
 economy and public services within the <u>Southampton Data Observatory</u>
- Health and Wellbeing Boards should develop a Health and Wellbeing Strategy paying due regard to the evidence set out in the JSNA.
- The Southampton Health and Wellbeing Strategy is monitored using a key set of performance indicators (KPIs). These can be accessed via a regularly refreshed <u>Power BI dashboard</u>. They are also available to view (along with commentary) within this slide pack.





What does the JSNA tell us about Health & Wellbeing in Southampton?





Demography

Current population

Southampton had an estimated resident population of **263,769** in 2022, of which...

134,578 (51.0%) were **male** and **129,191** (49.0%) were **female**

Population age groups

Aged 0-15 – **45,717** (17.3%) Aged 16-64 – **180,284** (68.3%) Aged 65 and over – **37,768** (14.3%)


Population

Dashboard

Aged 16-24 - 48,818 (18.5%)

Compared to 10.5% nationally – Southampton has a very young population

Data source: Hampshire County Council, Small Area Population Forecasts(SAPF) 2022 base Percentage of population by sex for Southampton 2022




Population for Southampton 2022				
Age group	Female	Male	Total	
Aged 00-04	6,926	7,221	14,147	
Aged 05-09	7,149	7,521	14,670	
Aged 10-14	6,949	7,405	14,354	
Aged 15-19	10,322	10,535	20,857	
Aged 20-24	14,643	15,864	30,507	
Aged 25-29	10,314	12,088	22,402	
Aged 30-34	9,690	11,388	21,078	
Aged 35-39	8,201	9,481	17,682	
Aged 40-44	7,360	7,985	15,345	
Aged 45-49	6,793	7,251	14,044	
Aged 50-54	7,036	7,180	14,216	
Aged 55-59	7,189	7,261	14,450	
Aged 60-64	6,192	6,057	12,249	
Aged 65-69	5,093	4,796	9,889	
Aged 70-74	4,781	4,406	9,187	
Aged 75-79	4,089	3,644	7,733	
Aged 80-84	2,905	2,350	5,255	
Aged 85-89	2,022	1,363	3,385	
Aged 90+	1,537	782	2,319	
Total	129,191	134,578	263,769	

southampton dataobservatory



Births




Births data details the **mothers birth region**, understanding this, along with births rates and changes in migration helps with **maternity service** and **school pupil place planning**.


In Southampton, the percentage of **mothers born outside the UK** is **increasing**.


The **birth rate** in Southampton remains **significantly higher** than England, although both are **falling** over time

Local rates are **falling faster** than nationally

In the **20% most deprived** areas, birth rates (12.4 per 1k) are **1.6x higher** than in the 20% least deprived (7.6 per 1k)







Proportion of total live births by mothers birth region in Southampton, 2015 to 2021

A **public health concern** is babies being born of **low birth weight** (under 2.5kg).

southampton

dataobservatory

In 2020, 3.8% of births were of low birth weight; significantly higher than England.

Bevois Ward was found to have the highest percentage of low birthweight babies. Local analysis shows Bevois has a higher concentration of Asian mothers who are more likely to have lower birth weight babies compared to the UK average. This reflects <u>published literature</u> where analysis confirms lower birth weight in second generation South Asian babies.


In **2018-20**, the percentage of low weight births in the **20% most deprived** areas (3.5%) was **3.2x higher** than in the **20% least deprived** (1.1%)

Source: Office for National Statistics



Population projections

southampton dataobservatory



Forecasts show a drop in residents aged under 16 (-1.8%), whilst the biggest increase is for those aged 65+ (+18.7%) between 2022 and 2029.

This is even greater for the 80+ age group, which is forecast to increase by +29.1%,

This ageing population will provide a future challenge and likely increase demand for health and social care services



southampton dataobservatory


Life expectancy and mortality

Life expectancy and mortality (southampton.gov.uk)

Life Expectancy

southampton dataobservatory

Poor health and **premature mortality** are **intertwined**. **Understanding** how long people are expected to live for (life expectancy), and how this **compares** locally with national average and comparator areas is an important measure of health.



Source: Office for Health Improvement and Disparities (OHID)

2019-21 national data expected November 2023


Southampton as a whole and at neighbourhood level has some of the **lowest life expectancies** in our **wider area**. It is difficult to see this on a map because of different population densities. In denser areas the detail is lost compared to more population sparse areas.

The next slide shows **cartograms** which are **maps reformatted** so the neighbourhoods of around 8,000 people cover the **same amount of diagram** space (regardless of land area covered)

In **2018-2020**, male life expectancy was **78.3** years in Southampton; significantly lower than England (**79.6** years) and ranking **5**th worst among comparators.

For **females** it was **82.5 years**; again **significantly lower** than that for England of **83.1** years and ranking **7**th **worst** among **comparators.**


Male and female life expectancies have followed national trends until for 2010-12 for males and 2014-16 for females; Southampton's rates have started to decrease whilst those for England have plateaued, which has widened the gap




2016-20 Life Expectancy – Cartograms (MSOA - 7.5k to 8k neighbourhoods) dataobservatory

Life expectancy at birth (females), Hampshire


and the Isle of Wight MSOAs: 2016 to 2020





34 neighbourhoods have **low male life expectancy** between 64.7 and 77.4 years

13 of these (38%) are in Southampton







Life Expectancy at birth - PCN level

southampton dataobservatory





Life expectancy at birth (Females) - Southampton Primary Care Networks: 2019 to 2021 (pooled)



Sources: NHS Digital Civil Registration Deaths Extract, GP Registration data

For both males and females, patients in the North PCN are expected to live longer than patients registered to other Southampton GPs, living 83.2 years and 85.1 years respectively.

Males living in the North PCN expect to live significantly longer than the Southampton male average

Males in Central PCN are expected to live the shortest time, for 79.3 years, just under 4 years less than those in North PCN.

Females in West PCN are expected to live the shortest time by sex, 82.6 years, 2.5 years less than those in North PCN




Life Expectancy at 65 years - PCN level

Life expectancy at 65 years (Males) - Southampton Primary Care

Networks: 2019 to 2021 (pooled)

southampton dataobservatory



Life expectancy at 65 years (Females) - Southampton Primary Care Networks: 2019 to 2021 (pooled)

Sources: NHS Digital Civil Registration Deaths Extract, GP Registration data

Sources: NHS Digital Civil Registration Deaths Extract, GP Registration data

After reaching age **65 years**, for both **males** and **females**, patients in the **North PCN** are expected to live the longest, living **21.3 years** and **23.1 years** respectively. **Bitterne PCN females** are also expected to lived **23.1 years** after turning 65 years old.


North PCN males and Living Well PCN males are expected to live significantly longer than the average male Southampton GP registered patients.


Males in West PCN are expected to live the shortest time after 65 years, for 17.7 years, 3.6 years less than those in North PCN. Females also in West PCN are expected to live the shortest time by sex, 20.6 years, 2.6 years less than those in North PCN




Life Expectancy at deprivation level

southampton dataobservatory





Life expectancy at birth (Females) - Inequalities Trend - Most Vs Least Deprived IMD England Quintiles (IMD 2019): 2014-16 to 2019-21 (pooled)



Sources: NHS Digital Civil Registration Deaths Extracts, ONS Mid-Year Population Estimates and IMD (2019)

Life expectancy at birth for males in the most deprived 20% has remained fairly constant, decreasing by 4 months between 2014-16 and 2019-21. In the least deprived 20% life expectancy has decreased for males by 11 months.

For females, life expectancy has decreased by just over a year for those in the most and least deprived 20% between 2014-16 and 2019-21.


The more recent decrease for those in the least deprived between 2018-20 and 2019-21 will be impacted by COVID-19. **Older affluent** people, who aged into **frailty**, were **more likely** to have **poorer outcomes** such as **pandemic related deaths**. Another **antecedent** was **multiple long-term conditions**, **more prevalent** in the **most deprived 20%**, these may both be factors in the **recent decrease** in the **life expectancy gap**.

Gap in life expectancy

southampton dataobservatory

Breakdown of the life expectancy gap between the most and least deprived quintiles of Southampton by cause of death, 2020 to 2021 (Provisional)

Percentage contribution (%)



Source: Office for Health Improvement and Disparities based on ONS death registration data (provisional for 2021) and 2020 mid-year population estimates, and Department for Levelling Up, Housing and Communities Index of Multiple Deprivation, 2019.

COVID-19 contributed **4.3%** to the gap in **male life expectancy** and **4.2%** to the gap for **female life expectancy**.

The chart shows the relative contribution that **nine broad causes of death** have on the **gap between life expectancy** for **Southampton** the **most deprived** and **least deprived quintiles of Southampton** 2020 to 2021 period.

Males

Circulatory (20.5%) **cancer** (19.7%) and **respiratory** (19.2%) **deaths** are the largest groups contributing to the **gap** in **male life expectancy** the **most deprived** and **least deprived quintiles of Southampton**. A deeper data dive shows the two largest causes are **chronic lower respiratory disease** followed by **heart disease**.

Females

Circulatory diseases (24.6%) is also the largest group contributing to the **gap** in **female life expectancy** between the **most deprived** and **least deprived quintiles of Southampton** with **respiratory diseases** (23.6%), other causes (14.9%) and external causes (10.5%), **cancer** was only **3.5%** for **females**, unlike for **males** where it was over **5 times higher**.


More detailed analysis shows the single largest causes of the gap in female life expectancy is chronic lower respiratory diseases followed by other causes and lung cancer.



Life Expectancy and healthy life expectancy

southampton dataobservatory

Data from the Annual Population Survey, calculates **healthy life expectancy**, which is a measure of how long people live in good health. **Life expectancy** MINUS **healthy life expectancy = Years in poor health** which is illustrated below



Females in the city may **live longer** than **males** (82.3 years versus 78.0 years) but they live in **poorer health** for **longer** (16.9 years versus 15.6 years).


2019-21 data is provisional, official statistics expected to be published November 2023



Life expectancy and healthy life expectancy


southampton dataobservatory

Life expectancy compared with healthy life expectancy for MALES in Southampton, by England deprivation quintiles, 2019-21*



Source: NHS England and ONS using ONS Silcocks method for Life Expectancy and ONS Sullivan method for Healthy Life Expectancy, *provisional data

Life expectancy compared with healthy life expectancy for FEMALES in Southampton, by England deprivation quintiles, 2019-21*



Source: NHS England and ONS using ONS Silcocks method for Life Expectancy and ONS Sullivan method for Healthy Life Expectancy , * provisional data

Females in the city may **live longer** than **males** but they live in **poorer health** for **longer** which ever deprivation quintile they live in.

Looking at **life expectancy versus healthy life expectancy**, in the **most deprived 20% England quintile**s (used by Core20+5 analysis), **males** live on average for **18.4 years** in **ill health** however females live for **19.2** years in ill health. Both males and females in the **most deprived quintile** live a **quarter (24%)** of their **shorter** lives in ill health. **Males** and **females** in the **least deprived** quintile live a **seventh (15%)** of their lives in **ill health**.

2019-21 life expectancy data expected to be published November 2023


Mortality – Years of life lost in Southampton 2019 to 2021

southampton dataobservatory

Some causes of death occur earlier in the life-course than others and therefore have a larger impact. We can measure this through calculating Years of Life Lost


Years of life lost are calculated by summing the number of years between the age at death and 75 years of age. This helps illustrate which causes of death have the greatest impact on life expectancy and young people

Analysis of these **trends**, **patterns** and **comparisons** helps us understand **priorities** for **health** and **wellbeing**



Mortality – Underlying causes of deaths Southampton 2021

southampton dataobservatory



Analysis of the **trends**, **patterns** and **comparisons** for cause of death helps us understand **priorities** for **health** and **wellbeing**

Comparing proportions of **deaths by cause** with proportions of **years of life lost by cause** shows which groups impact younger people disproportionately:

External causes account for 4.1% of deaths in 2021 but 14.5% of years of life lost.

Suicide and injury undetermined are the largest part of this accounting for 3.1% of deaths and 7.4% of years of life lost

Drug related deaths account for 0.9% of deaths in 2021 and 5.3% of years of life lost

Liver disease (incl. cirrhosis) is the underlying cause for 1.9% of deaths and 6.3% of years lost





Health conditions


Health conditions (southampton.gov.uk)



Chronic/Long-term conditions (LTCs)

southampton dataobservatory

- An ageing population compounds the prevalence of chronic/long-term conditions as people tend to develop more long-term or chronic conditions as they grow older
- Age analysis shows multi-morbidity increases with age, by 40-44 over half of residents have at least one chronic/long-term condition and by 80-84 over a third will have at least six long term conditions
- Analysis of snap shots from 2021 GP patient data shows more diagnoses of multiple chronic/ long-term conditions earlier in their life course than in 2017




Source: Sollis Clarity Health Analytics (ACG version 11.1/11.2) February 2021

Number of chronic conditions by age band Southampton patients February 2021



Long term conditions – Multi-morbidity and inequalities

southampton dataobservatory



- Multi-morbidity increases with age, BUT it also appears to be occurring earlier in life.....particularly for those in the most deprived areas
- Comparing **20% most deprived** and **20% least deprived** areas for prevalence of 3+ and 5+ chronic conditions:
 - differences appear at an early age and gradually narrow (relatively) as the population ages
 - differences begin to appear in the early 20's and peak between the mid-30's and mid-40's, where
 prevalence is more than double for those with 3+ conditions and more than triple for 5+ conditions



Leading causes and risk factors of disability



Understanding the leading causes and risks contributing to disability helps inform health and wellbeing action

Causes	Southampton	Portsmouth	Hampshire	Isle of Wight	England
Low back pain	1	1	1	1	1
Diabetes	2	2	2	2	2
Depressive disorders	3	3	3	3	3
Headache disorders	4	4	4	4	4
Neck pain	5	5	6	6	5
Gynecological diseases	6	6	9	13	9
Other musculoskeletal	7	7	8	10	7
Anxiety disorders	8	8	11	12	10
Age-related hearing loss	9	9	5	5	6
Asthma	10	11	12	8	14

Top 10 Causes attributed to Years Lived with Disability (YLDs)

Low **back pain** and **diabetes** are the **two** leading **causes** of disability across the local area and nationally

Risks Factors	Southampton	Portsmouth	Hampshire	Isle of Wight	England
High body-mass index	1	1	1	2	1
High fasting plasma glucose	2	2	2	1	2
Smoking	3	3	3	3	3
Alcohol use	4	4	4	4	4
Drug use	5	5	5	5	5
Occupational ergonomic	6	6	8	8	7
High blood pressure	7	7	6	6	6
Low bone mineral density	8	8	7	7	8
High processed meat	9	9	9	9	9
Particulate matter	10	10	10	12	10

Top 10 Risk Factors attributed to Years Lived with Disability (YLDs)

High body mass index and high fasting plasma glucose are the two leading risk factors causing disability across local area and nationally



Leading causes and risk factors of disability

southampton dataobservatory

Top ten conditions causing greatest disease burden

The top ten causes shown in the table below account for 37.3% of total DALYs in the selected area (or closest region if an ICB has been selected or parent county if a district has been selected).

Top ten conditions causing greatest disease burden (Disability-Adjusted Life Years): Southampton

Cause Name	Percentage of total DALYs in selected area (%)		
Ischemic heart disease	6.45		
Low back pain	4.96		
Chronic obstructive pulmonary disease	4.42		
Tracheal, bronchus, and lung cancer	3.86		
Diabetes mellitus	3.64		
Stroke	3.10		
Depressive disorders	3.09		
Headache disorders	2.85		
Falls	2.65		
Drug use disorders	2.29		

Top 10 conditions causing greatest burden measured in disabilityadjusted life years (DALYs)

Ischemic heart disease is the most common condition causing greatest burden with **Stroke** placed **6**th

COPD is the condition with the **3**rd greatest burden and **Diabetes** being the **5**th

Majority of causes have smoking as an upstream factor


Source: Global Burden of Disease 2019. Institute for Health Metrics and Evaluation (IHME). GBD Compare Data Visualization. Seattle, WA: IHME, University of Washington, 2020. Available from http://vizhub.healthdata.org/gbd-compare. (Accessed 06/09/2022)

Note: GBD 2019 data are only available for area geographies as at 2019. As such, no data are available for the 2021 geographies of North Northamptonshire and West Northamptonshire. GBD values displayed for these areas are for the former geography of Northamptonshire. Likewise, no data are available for the 2021 geography of Bournemouth, Christchurch and Poole. GBD values displayed for this area are for the former geography of Bournemouth.



Chronic/Long-term conditions (LTCs)


southampton dataobservatory




A <u>data pack</u> mapping the GP **diagnosed prevalence** of **18 common chronic/long-term conditions,** and 3-5+ **multiple conditions** across the city is available. This also includes **modelled forecasts of disease prevalence** by **age** and **locality** for these **conditions** in the **future.**

The top **FOUR** diagnosed conditions of Southampton registered patients are **hypertension**, **frailty**, **asthma** and **diabetes**.

Additional logistic modelling using the **GP data** <u>and</u> **Health Survey for England data** estimated **5,600** residents need for **help** with 5 or more **activities of daily living** in 2022, which is expected to increase by **+14%** to 6,400 **by 2028**









Childhood obesity and the food environment

Healthy weight (southampton.gov.uk)




- The leading cause of disability is a high body mass index (slide 22).
- Obesity in children is a risk factor for obesity in adulthood, which is a leading cause in a vast range of conditions*. (*Conditions such as asthma and other respiratory problems, eating disorders, mental health disorders and psychosocial risks, cardiovascular diseases, Type 2 diabetes, musculoskeletal problems, sleep apnoea etc.)
- Before the pandemic, a Scrutiny enquiry recommendation on childhood obesity was that analysis was conducted on childhood obesity and the food environment. Analysis on <u>childhood obesity</u> and the <u>food environment</u> was provided for a Task & Finish Group, available on the JSNA in the resources section of the <u>Healthy weight JSNA topic page</u>.
- In Southampton the level of obesity among year R children has remained stable and similar to the national average, whereas rates in year 6 children have increased overtime and have become worse than England.
- During the **COVID-19** pandemic, data was collected from a **representative sample (2020/21)**. Reception Year data for this period showed a **significantly higher increase** for obesity (17.1%) and excess weight (32.7%) prevalence locally and nationally compared to the previous four years.
- The Year 6 2020/21 sample for Southampton was **too small** to make **robust** statistical comparisons. However, the prevalence for **Year 6 obesity** (26%) and **excess weight** (41%) **mirrored** the **national** figures and **increasing prevalence** in the trend data follows the **national direction** of travel.
- The data also showed the gap in obesity prevalence between children in the most and least deprived parts of Southampton has widened. Linked analysis showed 7 out of 10 overweight Year 6 children and 4 out of 10 obese Year 6 children were of a healthy weight previously in Reception year.



Reception year obesity trend data

southampton dataobservatory



Year R Obesity and Excess Weight - Southampton and England trend: 2016/17 to 2020/21

Source: NHS Digital NCMP Enhanced data sets 2016/17 to 2020/21 with 95% Confidence Intervals (Wilson)

2021/22 England - Year R:Obese 10.1%Excess Weight 22.3%Southampton - Year R:Obese 10.9%Excess Weight 22.1%

Between **2016/17** and **2019/20** the level of childhood **obesity** and **excess weight** for **reception year children** locally and nationally largely **remained** at statistically **similar levels***.

*(Except for in 2017/18 Southampton had a significantly higher level than the national average for Year R obesity)

However, **2020/21** shows a **significantly higher increase** for obesity and excess weight prevalence in **year R locally** and **nationally** compared to the previous four years.

The latest data for **2021/22** shows Southampton had similar rates of childhood obesity and excess weight when compared to England.

Year 6 obesity trend data

southampton dataobservatory



Between 2016/17 and 2019/20 the level of childhood obesity and excess weight for Year 6 children has been higher locally than nationally and significantly higher in 2017/18 and 2019/20, Local level were also significantly higher than England for obesity in 2018/19.

The **gaps** between **Southampton** and **England** percentages of obesity and excess weight has **narrowed** in the **last two years**.

The latest data for **2021/22** shows **Southampton** had **higher** but not significantly rates of childhood **obesity** and **excess weight** when compared to **England**.

2021/22 England - Year 6: Southampton - Year 6: Obese 24.9% Obese 23.4%

Excess Weight 39.5% Excess Weight 37.8%


Linked analysis – changes in children's weight status in Southampton dataobservatory

Analysis linking **individual children's weight** recorded in **Year R** with that recorded for **Year 6,** gives **insight** into childhood obesity patterns in our city

Analysis showed although Year R obesity is a predictive factor for obesity in Year 6, interventions targeted only at this cohort only have the potential to reduce Year 6 obesity by one third.



The **majority** of **overweight** children in **year 6** had been **healthy weight** in **reception**, whilst just below a fifth had remained overweight and a further 9% had been obese





Over two-thirds (67%) of obese children had not been obese in reception, in fact the biggest proportion was for those who had been healthy weight (41%)




The Food Environment


southampton dataobservatory

Food environment impacts on childhood obesity

Fast food outlet data highlighted the **majority of residents** live with a **5-10 minute** drive or a **1km walk** of a **fast food** outlet

Almost all residents are within a mile of a fast food outlet, 7 out of 10 schools are within 400m of a fast food outlet, with closer proximities in the city centre and deprived areas.





Access to **supermarkets** with **larger floor spaces** (2,800+ m²) holding **more range** and more likely to include **budget brands** is **further** away from people in the **East** of the city and **Bassett** and **Swaythling**.

People in **deprived** areas are **less likely** to order groceries **online**

The full food environment analysis is on the Data Observatory





Diabetes

Diabetes (southampton.gov.uk)




- Diabetes is the second largest contributor to years of healthy life lost due to disability (YLDs) in Southampton and high fasting plasma glucose is the third biggest risk factor for deaths in Southampton, increasing the risk of cancer, cardiovascular diseases and neurological diseases (GBD 2019).
- Prevalence of diagnosed diabetes in Southampton (6.2%) is lower than the England average (7.3%) and is lower than most of its comparators (possibly due to its relatively young population). However, prevalence has been increasing in Southampton (+14.8% increase since 2012/13 but not as steeply as England +20.7%).
- **Despite** the **lower prevalence**, those people who do have **diabetes** in Southampton have some of the **worst outcomes in England**.
- Southampton's ratio of diabetic complications, rates of diabetic eye conditions and rates of minor diabetic lower limb amputations are all significantly higher than the England average and are the highest amongst Southampton's comparators.
- Southampton has an **ageing population**, this alone would result in nearly **1,500 additional cases** of diabetes in Southampton by **2028**. If Southampton's **prevalence rate continues to grow as well**, this increase could be greater than **+10,000 more cases by 2028**.



Mapping diabetes in Southampton


southampton dataobservatory



By PCN, Bitterne PCN has the highest prevalence (7.6%)

- The 20% neighbourhoods with the lowest prevalence are mainly in the centre of the city.
- The 5 LSOAs with the highest prevalence of diabetes are spread across Southampton. They are all located on or near the outer edge of the city.

Percentage of patients aged over 16 years with diabetes mellitus, Southampton PCNs: 2020/21







Respiratory

Respiratory (southampton.gov.uk)



Respiratory disease in Southampton (1)

southampton dataobservatory

- Chronic respiratory diseases ranked 3rd highest cause of Southampton deaths in all ages with a rate of 67.1 per 100,000. This is the same rank as it was in 1990. Respiratory infections and tuberculosis are ranked 5th highest cause of death for all ages in 2019 with a rate of 47.2 per 100,000 (GBD 2019).
 Asthma was ranked 10th highest for years of life lived with disability (YLD) with a rate of 382.6 YLD per 100,000 for all ages, a decrease of 42.1% since 1990. COPD was ranked 12th highest for years of life lived with disability (YLD) with a rate of 331.8 YLD per 100,000 (GBD 2019).
- Smoking and second-hand smoke is one of the biggest risks for respiratory diseases. Around 1 in 6 people (16.8%) in Southampton smoke. Higher when compared with 13.9% in England and 10.0% in Hampshire. More males smoke than females and people who smoke are more likely to be between the ages of 25 and 54
- Respiratory deaths contribute 19.2% of the gap in male life expectancy between the most and least deprived quintiles (2020-2021). On closer inspection, the two largest causes are chronic lower respiratory disease followed by heart disease (OHID Segment tool)
- For females, respiratory diseases contribute 23.6% of the gap in life expectancy between the most and least deprived quintiles, the 2nd highest group. More detailed analysis shows the single largest cause of the gap in female life expectancy is chronic lower respiratory diseases followed by other and lung cancer (OHID Segment tool).

- Rates of respiratory disease hospital admissions are higher for residents in the west of the city, especially who live in Redbridge and Coxford. Inequalities by deprivation shows admission rates for respiratory disease are 2.9x higher (and significantly so) for those in the 20% most deprived (England quintiles/Core 20+5) compared to the least. Under 75 years respiratory mortality rates are 2.5x higher for those in the 20% most deprived (England quintiles/Core 20+5) compared to the least. Under 75 years respiratory mortality rates are 2.5x higher for those in the 20% most deprived (England quintiles/Core 20+5) with the highest rates in Swaythling and Bargate
- Asthma prevalence rates are 1.2x higher for those in the 20% most deprived (England quintiles/Core 20+5) with the highest rates also in the west around Redbridge and Coxford. Asthma is more prevalent in those aged 60 to 84 years. Asthma correlates with current and ex-smokers. Asthma under 18 admissions rates are higher in the top two most deprived quintiles, in particular for 0–9 years, rates highest in the most deprived 20%
- **COPD prevalence** is **higher** on the **wings** and **edges** of the city, **highest rates** are found in **the west** city neighbourhoods with in **Coxford** and **Redbridge** and also for those aged **75 to 89 years.**
- COPD inequalities analysis shows by England quintiles COPD prevalence 1.2x higher, COPD admissions 3.4x higher and COPD mortality 2.6x higher for those in the most deprived 20% compared to the least





Cardiovascular

Cardiovascular disease (CVD) (southampton.gov.uk)



- Cardiovascular disease is the second highest ranking disease in Southampton for deaths and disability adjusted life years (DALYs) for all ages and rises to the highest rank for those aged 70 and over (GBD 2019).
- Circulatory diseases, including stroke, heart disease and CVD deaths contribute 20.5% of the gap in Southampton male life expectancy between the most and least deprived quintiles and is the largest group identified. For females, circulatory diseases contribute 24.6% of the gap in life expectancy between the most and least deprived quintiles, also the largest group identified (OHID Segment tool)
- Hypertension is estimated to be present in a third of the adult population. In Southampton, the known prevalence for 2021/22 is 10.8% or 32,550 patients. Other estimates by ONS suggest for every 7 adults diagnosed with hypertension there another 3 adults who are undiagnosed.
- NHS Health checks can identify help hypertension and early signs of stroke, kidney disease, heart disease, type 2 diabetes or dementia. The pandemic affected Health Checks and in Southampton, 2.8% of the targeted 20% eligible population (14.0%) had a NHS Health Check in 2021/22.



- Emergency hospital admissions for cardiovascular disease has highest rates for Bevois and then two wards in the west of the city; Coxford and Redbridge, the rate is 1.8x or 80% higher and for cardiovascular mortality 3.6x higher in the most deprived 20% of the city compared to the 20% least deprived
- Coronary heart disease emergency hospital admissions are 3.9x in most deprived 20% of the city compared to the 20% least deprived. The highest rates by ward are for people living in Redbridge (west Southampton), followed by Bitterne (now called Thornhill) and Bevois, all areas with high deprivation. Southampton has had higher coronary heart disease mortality rates than England since 2001-2003. At PCN level, Central PCN has the highest mortality rate compared to Southampton PCN average, followed by West PCN then Woolston and Townhill PCN
- Stroke prevalence in Southampton has been significantly lower than the England and more likely for those in the least deprived than the most deprived, perhaps occurring in affluent residents more likely to live longer when stroke risk is greater






Wider determinants and inequalities



Deprivation



southampton dataobservatory


- Southampton is ranked 55th (previously 54th) most deprived of 317 local authorities
- Around 12% of Southampton's population live in neighbourhoods within the 10% most deprived nationally (18% for the under 18 population)
- Southampton is ranked 3rd worst in the country for crime deprivation and is in the worst 20% of local authorities for FIVE other deprivation domains
- Grouping indicators by the deprivation levels (people experience living in these neighbourhoods) helps us explore inequalities within the city

\bigcirc

Inequalities – Children and Young People

southampton dataobservatory

Comparing outcomes for children and young people in the most deprived 20% of Southampton to the least deprived 20% illustrate the inequality gap in the city.....







Economic Needs Assessment

Economic assessment (southampton.gov.uk)



Southampton Wider Determinants Headlines (Economy focussed)

Analysis conducted on Southampton's economy contributes to our understanding of a number of **wider determinants of health and wellbeing**

southampton dataobservatory

The **<u>Economic Needs Assessment</u>** explores a **whole range** of areas that affects the **inequality gap** and also helps forecast the impact of areas of concern, for example the **cost of living**






Productivity and Growth – GVA (B)

southampton dataobservatory

Southampton's Economy in 2021



- Gross Value Added (GVA) is a key economic indicator
- Southampton represents 2.2% of South East economy
- The Southampton economy declined by -13.5% between 2019 and 2020, followed by an increase of +5.0% in 2021
- Since 2019, this represents a decline of -£681 million
- The England (+2.3%) and South East (+1.8%) economies have grown over the last two years, whilst the Southampton economy has declined since 2019 (-9.1%)
- Additionally, this is the greatest decline among comparators; the majority (8 out of 10) having experienced growth
- Overall this suggests that the economic impact of the COVID-19 pandemic was greater locally




GVA (B) Million £ at current basic prices- Southampton: 1998 to 2021

Change since 2019:



Source: Office for National Statistics - GVA. GVA rounded to nearest hundred million.



+1.8%

England


Source: Office for National Statistics - GVA



Claimant Count


southampton dataobservatory

Claimants as a proportion of residents aged 16-64 (Total) -Southampton and ONS comparators: March-2023




Claimants as a proportion of residents aged 16-64 (Total) - Southampton, England, South East: April-2021 to March-2023





- Locally and nationally the number of adults claiming out of work benefits has significantly decreased over the last two years, given the end of COVID-19 restrictions
- 4.2% (7,060) of the working aged population in Southampton were claiming out of work benefits in March 2023; a decline of -4,940 (-41%) since April 2021 (7.1%)
- Although, Southampton is yet to return to the pre-pandemic baseline (less than 3.5% in January to March 2020)
- Claimant count also appears to have slowly increased in recent months, possibly a result of recent financial pressures and economic uncertainty, therefore it will be important to monitor this trend



Source: Department for Work and Pensions




Claimant Count Inequalities

southampton dataobservatory


- The map below shows the latest claimant count (%) by Southampton neighbourhoods March 2023
- Higher claimant counts are seen across neighbourhoods in Thornhill, Woolston, Bevois and Redbridge wards, which is where some of the most deprived neighbourhoods in the city are located
- The chart below shows the inequality gap in the claimant count between the most and least deprived neighbourhoods over time, which has decreased from a peak percentage point gap of 8.1 in April 2021 to 5.0 in March 2023, whilst the inequality gap briefly returned to pre-pandemic levels (average 4.6 percentage point gap throughout 2019), it appears to be widening again

Change in the claimant count for the most and least deprived national deprivation quintiles in Southampton: April-2021 to March-2023



Inequality gap 20% most deprived 20% least deprived

Source: Department for Work and Pensions



Claimant count (total) as a percentage of the working age population by LSOA: March-2023



Inequalities – Workplace vs Resident

southampton dataobservatory

Workplace to Resident pay gap, Weekly pay - gross, Full Time Workers, (Total) - Southampton and ONS comparators: 2022

Leeds -f40Hampshire -f38 South East -f21 Sheffield -f16 Bournemouth Ch... -f11 Newcastle upon ... Isle of Wight -f5 England -£0 Liverpool £11 York Portsmouth f18 Coventry £21 Bristol, City of £24 Southampton £37 Plymouth f40-£40 -£20 £0 £20 £40 Pay gap £

Workplace to Resident pay gap, Weekly pay - gross, Full Time Workers - (Total) Southampton: 2012 to 2022

Percentage pay gap Workplace earnings Resident earnings



- WORKPLACE earnings are £37 (5.7%) more per week than RESIDENT earnings for full time workers in Southampton
- Whilst the inequality gap between workplace and resident earnings appears to have narrowed over the last two years, the gap is still the second largest among comparators
- High workplace earnings suggests that good skilled employment opportunities exist in the city. However, lower resident earnings suggests that commuters into the city have those high skilled jobs, which residents are not benefitting from



Inequalities – Male vs Female

Percentage pay gap Male earnings


southampton dataobservatory

Male to Female pay gap, Weekly pay - gross, Full Time Workers - (Workplace) Southampton: 2012 to 2022

Male to Female pay gap, Weekly pay - gross, Full Time Workers - (Resident) Southampton: 2012 to 2022

Percentage pay gap Male earnings







- There is also a pay gap between male and female pay in Southampton, with this gap also experienced nationally
- In 2021, the full time resident weekly gender pay gap was 14.6% (£99) in Southampton, this compares to a gap of £106 (15.3%) nationally
- The full time workplace gender weekly pay gap in Southampton was similar at 13.0% (£94) in 2022
- No evidence that gap is narrowing for both workplace and resident

Cost of Living – Earnings

southampton dataobservatory




Source: ONS - Annual Survey of Hours and Earnings & Consumer Price Inflation



Gross weekly pay for full time workers - workplace analysis: Southampton trend: 2008 to 2022 (projection)

Source: Office for National Statistics – Consumer Price Inflation & Annual Survey of Hours and Earnings



- The graphs on the left show projections for adjusted resident and worker earnings based on **April 2022 inflation** (120.0 CPI all items)
- Assuming unadjusted earnings do not change, a 'real' decline of -8.3% in earnings could be expected for both residents and workers in Southampton – an impact of inflation increasing during the cost of living crisis
- Unadjusted weekly earnings would need to increase by at least £54 for residents and £58 for workers to negate the impact of inflation in 2022;
- This would be an unprecedented increase in unadjusted earnings locally, therefore would expect wage growth in Southampton to fall behind inflation, resulting in a **decline in 'real' earnings in 2022**
- ONS estimates of weekly earnings from August 2022, also suggests that 'real' pay is declining as a result of inflation





COVID Impact Assessment



 Most aspects of health and wellbeing covered by the JSNA were impacted by the pandemic including those monitored against the Health and Wellbeing Strategy

southampton dataobservatory

- Further analysis of the direct and indirect impacts of the pandemic are included in the Covid-19 Impact Assessment, set out in three sections; Healthy People, Healthy Living and Healthy Places
- Many impacts are yet to be fully realised and the Covid-19 Impact Assessment is refreshed regularly as more data is made available and further understanding reached. Future impacts suggest this winter would have an impact on health and wellbeing inequalities in the community given the challenges of heating costs and the impact of the cost-of-living increase.
- The assessment showed **significant impact** of the **Covid-19 pandemic** on the **health of Southampton residents.** Analysis including looking at **inequalities**, showing there were **significant differences** in **cases** (in the first three waves) and **hospital admissions** when comparing those living in the 20% most deprived neighbourhoods with those living in the 20% least deprived - with **higher rates in the most deprived**
- There have been some **negative impacts** such as an **increase in mental health issues** but also some **positive impacts** such as **reduction in smoking, increased value of air quality** and **clean air**, and an increase in **physical activity**.
- Analysis incorporates national and local data including Southampton resident survey data

Covid-19 Impact Assessment



southampton dataobservatory

Other summary slides



Key facts

southampton dataobservatory

37,183

4th quintile

9778


20% least


deprived

Southampton has an estimated population of **263,767** residents, of which **134,575** (51.0%) are male and 129,192 (49.0%) are female (2022).

Southampton has a relatively young population compared to geographic neighbours with higher rates of deprivation, diversity and SC Rank 6: E01017245 IMD Score: 54.7 pre-existing disease. A shift towards an ageing population has been forecast for the city.

Deprivation is generally associated with poor health outcomes.





This map shows how deprivation is distributed across different neighbourhoods in the city with red areas experiencing much higher deprivation compared to blue areas.

20% most

deprived

2nd

quintile

3rd

quintile

Deprivation guintiles

50,000

The Index of Multiple Deprivation consists of 7 domains including income, employment, health and disability, education, crime, housing and living environment.

Southampton is ranked the 55th (previously 54th) most deprived out of 317 local authorities in England. 28% of Southampton's population live in neighbourhoods within the 20% most deprived nationally. Southampton is ranked 3rd worst in the country for crime deprivation and is in the worst 20% of local authorities for 5 other deprivation domains.